



Estrogen Receptor Luciferase Reporter MDA-MB-453 Stable Cell Line

Catalog Number SL-0087 (For Research Use Only)

Introduction

Estrogen receptor (ER) belongs to the nuclear receptor family and plays a widespread role in human physiology and the development or progression of numerous diseases. In response to estrogen stimulation, estrogen bound receptor in the nucleus dimerizes and binds to specific response elements known as estrogen response elements (EREs) located in the promoters of target genes and regulates their gene expression. Signosis has established the MDA-MB-453 ER luciferase reporter stable cell line, in which the ERE and reporter luciferase gene are consistently expressed in the cell line to facilitate the screening and study. This stable cell line can provide a sensitive, responsive, and rapid in vitro system to detect and measure substances with potential (anti-) estrogenic activity.

Product description

The cell line was established by transfection of ER luciferase reporter vector along with neomycin expression vector followed by neomycin selection. The G418 resistant clones were subsequently screened for 17 beta-estradiol induced luciferase activity. The clone with the highest fold induction (10 fold) was selected and expanded to produce this stable cell line.

Materials provided


One vial of 2×10^6 cells, at passage 4, in Freezing Media. **IMPORTANT:** store the frozen cells in liquid nitrogen until you are ready to thaw and propagate them.

Handling cells upon arrival



It is strongly recommended that you propagate the cells by following instructions as soon as possible upon arrival.**

IMPORTANT: It is imperative that an adequate number of frozen stocks be made from early passages as cells may undergo genotypic changes. Possible genetic instability in transfected cells may result in a decreased responsiveness over time in normal cell culture conditions.



Required Cell Culture Media

- **Complete Growth Media**
In 500mL of RPMI-1640, add 50mL FBS (10% final) and 5mL Penicillin/Streptomycin (1% final).
- **2x Freezing Media**
Add 10% DMSO (final) to Complete Growth Media and sterile filter. Make fresh each time.

Materials required but not provided

(Maybe substituted with a comparable third-party product)

Materials	Product number
RPMI-1640 Medium	Cytiva SH30027.01
Fetal Bovine Serum (FBS)	Cytiva SH30910.03
Penicillin/Streptomycin	Cytiva SV30010
Trypsin	Cytiva SH30042.02
Phosphate-buffered saline (PBS)	Cellgro P/N 21-040-CV
DMSO	Sigma P/N D8418
96-well white plate	Greiner Bio-One P/N 655098
Luciferase substrate	Signosis P/N LUC015
Cell lysis buffer	Signosis P/N LS-001
G418 (optional)	Invitrogen P/N 10131-027

Initial Culture Procedure

1. Quickly thaw cells in a 37°C water bath with careful agitation. Remove from the bath as soon as the vial is thawed.
2. Transfer cells to a 100mm² dish (or T-25cm² flask) containing 10ml of Complete Growth Media.
3. Gently rock the flask to ensure the cells are mixed well in the media. DO NOT PIPET.
4. Place the flask with cells in a humidified incubator at 37°C with 5% CO₂.
5. After cells adhere (wait at least 8 hours to overnight), replace media with fresh Complete Growth Media.

Subculture Procedure

1. After Cells have recovered and growing well subculture/passage cells when the density reaches 90-100% confluency, maintain and subculture the cells in Complete Growth Media.

Note: During the time that cells are not used for the experiment ideally, they can be maintained in Complete Growth Media with 50-100µg/ml of G418.

2. Carefully remove the culture media from cells by aspiration.
3. Rinse cells with PBS, being careful to not dislodge attached cells. Then remove PBS by aspiration.
4. Add 1-2 mL trypsin/Tris-EDTA solution.
5. Incubate with trypsin for 2-5 minutes (or until detached). Confirm detachment by observation under the microscope.
6. Add 5-10ml of pre-warmed Complete Growth Media and gently pipet up and down to break the clumps.
7. Passage cells in 1:3 to 1:5 ratio when they reach 90% confluency.

NOTE: Stable cell lines may exhibit a slower proliferation rate compared to parental cells. Do not seed cells at suboptimal density as this may hinder cell growth and division.

Preparing frozen stocks

This procedure is designed for 100mm²dish or T-75cm² flask. Scale volumes accordingly to other vessels.


1. When cells reach 90-100% confluency, freeze down cells.
2. Detach cells according to “Subculture Procedure.”
3. Transfer cells to a 15ml conical centrifuge tube and centrifuge at 250 x g (or 2,000 RPM) for 5 minutes to collect the cells into a pellet.
4. Carefully aspirate the media and resuspend cells in 0.5mL complete growth media.
5. Add 0.5mL of **2X Freezing Media** and gently resuspend by pipetting up and down.
6. Transfer 1mL of cells into a cryogenic vial.

7. Place the cryogenic vial in a freezing container (Nalgene # 5100-0001) and store it at -80°C freezer overnight.
8. Transfer cells to liquid nitrogen for long-term storage.

Assay procedure

The following procedure should be followed as a guideline. You will need to optimize the assay conditions based on your experimental setup.

1. The day before performing the assay, trypsinize the cells and seed each well of a **white clear-bottom** 96 well plate with 1-3 x 10⁴ cells in 100µl medium.
2. Incubate the plate in a humidified incubator at 37°C with 5% CO₂ overnight.
3. Add inducing reagent directly to each well and incubate for an appropriate time to produce maximal induction.
4. Remove the media by aspiration and add 100µl of PBS to each well.
5. Remove PBS by aspiration and add 20µl of 1x lysis buffer to each well (To prepare 1x lysis buffer, add one volume of 5x lysis buffer to four-volume of distilled water).
6. Incubate cells in lysis buffer for 15 minutes at room temperature.
7. Add 100µl of luciferase substrate to each well and gently pipette up and down.
8. Immediately read the plate in a luminometer with a sensitivity of 3×10⁻²¹ moles luciferase.



ER Luciferase Reporter MDA-MB-453 cells were treated for 16 hours at varying concentrations of 17β-Estradiol.

Signosis Luciferase Reporter Stable Cell Lines

For a complete list of cell lines please visit our website at <http://www.signosisinc.com/category/cell-based-assays>

Transcription Factors	Pathway	Cell Line	Catalog Number
NFkB	NFkB	Hela; human cervical cancer	SL-0001
NFkB	NFkB	NIH/3T3; mouse fibroblast	SL-0006
NFkB	NFkB	HEK293; human embryonic kidney	SL-0012
NFkB	NFkB	MCF-7; human breast cancer	SL-0013
NFkB	NFkB	A549; human lung cancer	SL-0014
NFkB	NFkB	HepG2; human liver cancer	SL-0017
NFkB	NFkB	Neuro2a; mouse neuroblastoma	SL-0026
NFkB	NFkB	MEF; murine embryonic fibroblast	SL-0033
NFAT	Calcium Signaling	Jurkat T; human T lymphocyte	SL-0032
NFAT	Calcium Signaling	Hela; human cervical cancer	SL-0018
NFAT	Calcium Signaling	NIH/3T3; mouse fibroblast	SL-0029
p53	p53	Hela; human cervical cancer	SL-0011
p53	p53	RKO; human colon cancer	SL-0007
SMAD	TGFbeta	HepG2; human liver cancer	SL-0016
SMAD	TGFbeta	NIH/3T3; mouse fibroblast	SL-0030
NRF2	Antioxidant Response	MCF7; human breast cancer	SL-0010
STAT1	JAK-STAT	Hela; human cervical cancer	SL-0004
STAT3	JAK-STAT	Hela; human cervical cancer	SL-0003
HIF	Hypoxia Response	NIH/3T3; mouse fibroblast	SL-0005
HIF	Hypoxia Response	Hela; human cervical cancer	SL-0023
HIF	Hypoxia Response	Neuro2a; mouse neuroblastoma	SL-0027
ER	Estrogen Receptor Signaling	T47D; human breast cancer	SL-0002
AR	Androgen Receptor Signaling	MDA-MB-453; human breast cancer	SL-0008
GR	Glucocorticoid Receptor Signaling	MDA-MB-453; human breast cancer	SL-0009
GR	Glucocorticoid Receptor Signaling	Hela; human cervical cancer	SL-0021
AP-1	JNK, ERK, MAPK Signaling	Hela; human cervical cancer	SL-0019
CREB	cAMP, PKA, CaMK Signaling	HEK293; human embryonic kidney	SL-0020
CREB	cAMP, PKA, CaMK Signaling	NIH/3T3; mouse fibroblast	SL-0031
CHOP	Unfolded Protein Response, ER stress	Mia-Paca2; human pancreatic cancer	SL-0025
TCF/LEF	Wnt/b-catenin	HEK293; human embryonic kidney	SL-0015
TCF/LEF	Wnt/b-catenin	Hela; human cervical cancer	SL-0022
TCF/LEF	Wnt/b-catenin	CHO-K1; Chinese Hamster Ovary	SL-0028
ELK	MAPK Signaling	HEK293; human embryonic kidney	SL-0040
ELK	MAPK Signaling	Hela; human cervical cancer	SL-0041
IRF	Immune Response Pathway	HEK293; human embryonic kidney	SL-0035

** Signosis products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to this user manual or product information sheet and shipped directly by Signosis.